



Post-Operative and Home Care Instructions for Breast Augmentation Surgery

DIET: Start with clear liquids and toast or crackers. If those are well tolerated, progress to a regular diet.

DRIVING: No driving for 1 week after your procedure or while taking pain medicine.

ACTIVITY: You may walk and climb stairs immediately after surgery. After 4 weeks you may resume light activity such as brisk walking. During the first 4 weeks do not lift anything heavier than a gallon of milk. After 6 weeks you may resume more strenuous aerobic work and lifting activities as discussed with your physician.

WORK: Depending on your career and your rate of healing, you should be able to return to work within 1 to 2 weeks of surgery. Your surgeon will be able to give you a better estimate depending on your physical and professional profile.

WOUND CARE: Your dressing may get wet in the shower. Try to direct the shower to your back, but it is fine if the dressing gets wet. Pat dry after your shower.

BATHING: You may shower and wash your hair 48 hours after surgery. Shower only for the first 30 days post-operative. Try to keep the shower directed at your back and not at your incision or dressing. It is okay if the dressing gets wet in the shower. Pat it dry after your shower. For the first 4-6 weeks after surgery, do not submerge in a bathtub, pool, or hot tub.

MEDICATIONS: Ask your surgeon when you may resume your blood thinning medications. All other prescription medications may be resumed the following day as long as you are tolerating fluids/liquids. While you are taking pain medicine, you are encouraged to follow a high fiber diet or take a stool softener such as Colace (available over the counter), as pain medications tend to cause constipation.

SMOKING AND ALCOHOL: Do not smoke for 4 weeks after surgery as it impedes wound healing and can lead to serious wound complications. Alcohol consumption is dangerous while taking pain medicine. It has a tendency to worsen bleeding.

POST-OPERATIVE APPOINTMENT: Your first follow-up visit will be 1 week after surgery. Your surgeon will then see you at appropriate intervals after this to monitor your progress.

SPECIAL CONSIDERATIONS: Call your surgeon immediately if you experience any of the following: excessive pain, swelling, bleeding, redness at the incision site, or fever over 101° F.